

ASSOCIATION OF FV G1691A, FV H1299R, AND FII G20210A VARIATIONS WITH THROMBOSIS AND CORONARY ARTERY DISEASE (CAD): A POPULATION-BASED STUDY

ASOCIJACIJA VARIJACIJA FV G1691A, FV H1299R I FII G20210A SA TROMBOZOM I BOLESTIMA KORONARNIH ARTERIJA (CAD): STUDIJA NA POPULACIONOM NIVOU

Sevda Ünallı Özmen^{1,2}, Yeşim Özarda³, Aylin Köşeler⁴, Ramazan Sabırlı⁵,
Derya Sucu Kaynak⁶, Ibrahim Koç⁷

¹Department of Medical Biochemistry, Central Laboratory, City Hospital, Bursa, Turkey

²Department of Translational Medicine, Health Sciences Institute, Uluda University, Bursa, Turkey

³Department of Medical Biochemistry, Faculty of Medicine, Yeditepe University, Istanbul, Turkey

⁴Department of Biophysics, Faculty of Medicine, Pamukkale University, Denizli, Turkey

⁵Department of Emergency Medicine, Serfergazi State Hospital, Denizli, Turkey

⁶Department of Occupational Health and Safety Specialist, Head Office, Bursa, Turkey

⁷Department of Pulmonary Medicine Clinic, City Hospital, Bursa, Turkey

Summary

Background: Thrombosis and coronary artery disease (CAD) are complex disorders influenced by genetic factors. Specific gene variations, such as Factor V (FV) G1691A (Leiden), FV H1299R, and Prothrombin (FII) G20210A, have been implicated in thrombotic events and CAD. However, their precise role in CAD development remains controversial. This study investigated the prevalence and association of these gene variations with thrombosis and CAD in the Turkish population.

Methods: A case-control study included 406 healthy individuals and 64 CAD patients. Genotyping for FV G1691A, FV H1299R, and FII G20210A was performed using a strip assay. Fisher's exact test compared allele and genotype frequencies between the CAD and control groups.

Results: No significant differences were observed in genotype frequencies of FV G1691A, FV H1299R, and FII G20210A between the CAD and control groups ($p > 0.05$). Similarly, allele frequencies did not differ significantly between the two groups ($p > 0.05$).

Conclusions: The findings suggest that FV G1691A, FV H1299R, and FII G20210A variations may not play a significant role in the development of CAD in the Turkish pop-

Kratak sadržaj

Uvod: Tromboza i bolest koronarnih arterija (CAD) su složeni poremećaji na koje utiču genetski faktori. Specifične genetske varijacije, kao što su Faktor V (FV) G1691A (Leiden), FV H1299R i protrombin (FII) G20210A, su povezane sa trombotskim događajima i CAD-om. Međutim, njihova tačna uloga u razvoju CAD-a ostaje nejasna. Ova studija je istraživala prevalenciju i povezanost ovih genetskih varijacija sa trombozom i CAD-om u turskoj populaciji.

Metode: Studija slučaj-kontrola je obuhvatila 406 zdravih osoba i 64 pacijenta sa CAD-om. Genotipizacija za FV G1691A, FV H1299R i FII G20210A je izvedena korišćenjem strip testa. Fisherov egzaktni test je korišćen za poređenje učestalosti alela i genotipova između grupe sa CAD-om i kontrolne grupe.

Rezultati: Nisu uočene značajne razlike u učestalosti genotipova FV G1691A, FV H1299R i FII G20210A između grupe sa CAD-om i kontrolne grupe ($p > 0,05$). Slično tome, učestalost alela takođe nije pokazala značajne razlike između ove dve grupe ($p > 0,05$).

Zaključak: Rezultati ukazuju na to da varijacije FV G1691A, FV H1299R i FII G20210A možda nemaju

Address for correspondence:

Dr. Sevda Ünallı Özmen
Department of Medical Biochemistry, Central Laboratory, City
Hospital, Bursa, Turkey
e-mail: ozmendr@hotmail.com

ulation studied. These results are consistent with the existing conflicting literature on the association between these gene variations and CAD. Further research with larger sample sizes and diverse populations is warranted to elucidate the role of these variations in CAD pathogenesis.

Keywords: thrombosis, coronary artery disease, factor V gene, prothrombin gene, gene variations, allele frequencies

Introduction

Thrombosis poses a significant global health challenge due to its high rates of mortality and morbidity. The development of thrombosis involves various factors, including environmental, clinical, and genetic influences. Similarly, the development of coronary artery disease (CAD) entails a multifaceted aetiology, primarily encompassing a combination of traditional risk factors such as type 2 diabetes, dyslipidemia, arterial hypertension, cigarette smoking, as well as genetic predisposition. Factor V (FV) and Prothrombin (FII) gene variations are commonly associated with thrombotic events. Still, the role of specific variants, such as factor V Leiden (FVL), prothrombin gene (PT G20210A), and methylenetetrahydrofolate reductase (MTHFR) C677T polymorphisms in coronary artery disease (CAD) development remains controversial.

Role of Factor V Gene Variations in Thrombotic Events

Familial thrombophilia, a common cause of inherited thrombosis, is often associated with Activated Protein C Resistance (APCR). APCR is primarily caused by a single variation in the Factor V (FV) gene, known as Factor V Leiden (FVL), located on chromosome 1. The FVL mutation is inherited in an autosomal dominant manner. This variation involves the substitution of guanine with adenine at nucleotide 1691 in the FV gene, replacing arginine with glutamine at the 506th amino acid position (Arg506Gln) of the FV protein. This variation occurs at the site of the first molecular cleavage of factor Va by Activated Protein C (APC). Due to this variation, thrombin is continued to form by the prothrombinase complex (1–4).

The number of FVL mutant alleles influences the clinical manifestation of FVL mutation-related thrombophilia. Certain circumstantial risk factors, such as central venous catheters, organ transplantation, contraceptive use, advancing age, hormone replacement therapy, surgery, and prolonged immobilization, such as in travel, can have an additive effect on thrombotic events (5). Heterozygote genotype frequencies of FVL variation range from 3% to 8% in the United States and Europe general populations. These frequencies vary significantly, with higher

značajnu ulogu u razvoju CAD-a kod ispitanika u turskoj populaciji. Ovi nalazi su u skladu sa postojećom kontradiktornom literaturom o povezanosti ovih genetskih varijacija i CAD-a. Neophodna su dalja istraživanja na većim uzorcima i raznovrsnijim populacijama kako bi se razjasnila uloga ovih varijacija u patogenezi CAD-a.

Ključne reči: tromboza, bolest koronarnih arterija, gen za faktor V, gen za protrombin, varijacije gena, frekvencija alela

frequencies observed in European populations and lower frequencies in African, indigenous Australian, and Asian populations. The homozygous genotype frequency of the FVL variation is approximately 1 in 5000 individuals (5).

In addition to the FVL variation, sequencing of exon 13 of the FV gene has revealed a 4070A-G variation (His1299Arg; H1299R), known as the HR2 haplotype. This variation is associated with several other polymorphisms (6). The HR2 haplotype encompasses more than 12 different polymorphisms throughout the FV gene (collectively known as HR2), with seven polymorphisms predicting amino acid changes in the FV protein and resulting in protein modifications (7, 8). Carriership of the FV H1299R allele is associated with mild APCR and interacts with the FVL variation, leading to a more severe APCR phenotype (9–11). However, the specific thrombotic event risk associated with the HR2 haplotype remains unclear (3, 12). Nevertheless, the presence of both the FV H1299R allele and the FVL variation has increased the risk of thrombotic events even further (9, 13).

Role of Prothrombin Gene Variation in Thrombotic Events

Following the Factor V Leiden (FVL) variation, the prothrombin or FII G20210A variation is the second most common genetic risk factor associated with thrombosis (14–16). The G20210A mutation is found in approximately 1–4% of healthy individuals, and its prevalence varies among populations. The highest prevalence is observed in Europe, while G20210A heterozygosity is extremely rare in Asian, African, and Native American populations. The prevalence of G20210A homozygosity is approximately one in 10,000 individuals (17–19).

The FII G20210A variation is a genetic polymorphism located in the 3'-untranslated region of the prothrombin (FII) gene. It is caused by a single base pair substitution, where guanine (G) is replaced by adenine (A) at position 20210. The inheritance of this variation is autosomal dominant and is associated with elevated plasma levels of prothrombin (FII) (17). Additionally, the FII G20210A variation has been implicated in the risk of arterial disease. Young women with the FII 20210A allele were reported to

have a fourfold higher risk of myocardial infarction, while men had a 1.5-fold higher risk (18).

Adult individuals carrying the G20210A heterozygote genotype have a two- to fourfold increased risk of thrombosis, while children with this genotype face an even higher three- to fourfold increased risk. Furthermore, individuals with G20210A homozygosity tend to experience thrombotic events more frequently and at a younger age. The FII variation has also been associated with an increased risk of preeclampsia and pregnancy loss (19, 20).

This case-control and prevalence study aims to determine the allele frequencies of the FV gene G1691A-H1299R and FII gene G20210A mutations as genetic risk factors for thrombosis and coronary artery disease in both the healthy population and patients with coronary artery disease in the South Marmara region of Western Turkey.

Materials and Methods

Ethical Issues, Study Population, and Blood Sampling

Ethical approval for the study was obtained from the ethics committee of Uluda University, ensuring compliance with ethical guidelines. All participants were provided with information about the study and their rights, and their informed consent was obtained following the principles outlined in the Helsinki Declaration.

The study population consisted of 406 healthy volunteers (179 males and 227 females) between the ages of 18 and 45 years from the Bursa region. These individuals were included to determine the genotype distributions of the Factor V (FV) and Prothrombin (FII) polymorphisms in the healthy population. In addition, a group of 64 patients (12 females and 52 males) aged between 18 and 45 years with documented coronary artery disease (CAD), confirmed by coronary artery angiography, were included in the study. For all participants, blood samples were collected using EDTA tubes (Vacutainer, Becton Dickinson, U.K.).

Genotyping

Genotyping for cardiovascular disease (CVD) in this study was performed using the CVD-StripAssay (ViennaLab Labordiagnostika GmbH, Austria). This assay is based on the reverse hybridization principle for mutation analysis, allowing for the simultaneous detection of Factor V (FV) G1691A-H1299R and Prothrombin (FII) G20210A mutations.

Variation analyses were conducted following the manufacturer's recommendations. DNA was extracted from anticoagulated blood using the Invisorb®

Spin Blood Mini Extraction Kit (Invitek, Germany) and stored at -20 °C for later use. Subsequently, relevant gene sequences were amplified and biotin-labelled in a single amplification reaction using the PCR amplification method.

In brief, PCR amplification was carried out in a final volume of 25 µL, consisting of 15 µL of pre-prepared PCR amplification mix, 4.8 µL of diluted buffer, 1U of Taq DNA Polymerase (Fermentas), and 5 µL of DNA. The thermal cycler (Applied Biosystems 2720 Thermal Cycler, USA) program included an initial step of 94 °C for 2 minutes, followed by 35 cycles of 94 °C for 15 seconds, 58 °C for 30 seconds, 72 °C for 30 seconds, and a final extension step of 72 °C for 3 minutes.

Following amplification, the products were selectively hybridized using an Autolipa automated incubator (ProfiBlot T48, TECAN, Switzerland) on a test strip that contained immobilized allele-specific (wild-type and mutant) oligonucleotide probes arranged as an array of parallel lines. Biotinylated sequences bound to the strip were detected using streptavidin-alkaline phosphatase and colour substrates. The genotype of each sample was determined by comparing the staining pattern of the processed Test strip with the Decoder table enclosed in the assay kit. Each polymorphic position displayed one of the following staining patterns: Normal, Heterozygous, or Homozygous mutant (21).

Genotyping for cardiovascular disease (CVD) was performed using the CVD-StripAssay based on the reverse hybridization principle. DNA was extracted from anticoagulated blood, and PCR amplification was conducted following the manufacturer's recommendations. The genotype of each sample was determined by comparing the staining pattern on the Test strip with the Decoder table provided in the assay kit.

Statistical analysis

The frequency of Factor V (FV) Gene G1691A (Leiden)-FV Gene H1299R and Prothrombin (FII) Gene G20210A genotypes and alleles were analyzed using the Statistical Package for the Social Sciences program (SPSS for Windows, version 17.0; SPSS, Chicago, IL).

Fisher's exact test was employed to compare allele and genotype frequencies between the two groups. A p-value of less than 0.05 was considered statistically significant for all analyses, indicating a significant difference between the compared groups.

Results

The findings regarding the genotype frequencies of FV G1691A are presented in *Table 1*. The results indicate no significant difference between the

Table I Distributions of mutations in healthy individuals and CAD Group.

		Control Group n=406	CAD Group n=64	OR (95% CI for Genotypes)	p-value
Polymorphisms	Genotypes	n (%)	n (%)		
Factor V G1691A(Leiden)	AA	1 (0.25)	1 (1.56)	1.45 (0.70–2.89)	0.170
	GA	34 (8.37)	7 (10.93)		
	GG	371 (91.38)	56 (87.5)	0.95 (0.86–1.05)	
Factor V Gene H1299R	GG	2 (0.49)	0	1.32 (0.68–2.59)	0.544
	AG	41 (10.10)	9 (14.07)		
	AA	363 (89.41)	55 (85.93)	0.96 (0.86–1.06)	
Prothrombin G20210A	AA	0 (0)	0	N/E	0.500
	GA	19 (4.68)	1 (1.56)	0.33 (0.04–2.45)	
	GG	387 (95.32)	63 (98.44)	1.33 (0.99–1.07)	

OR=Odds Ratio; p-values are derived from the Fisher Exact Test; N/E=Not estimated; CI: Confidence Interval

Table II Distributions of Alleles in healthy individuals and CAD group.

		Control Group n=812	CAD Group n=128	OR (95% CI for Alleles)	p-value
Polymorphisms	Alleles	n (%)	n (%)		
Factor V G1691A	G	776 (95.56)	119 (92.96)	0.97 (0.92–1.23)	0.187
	A	36 (4.44)	9 (7.04)	1.58 (0.78–3.21)	
Factor V Gene H1299R	A	767 (94.45)	119 (92.96)	0.98 (0.93–1.03)	0.538
	G	45 (5.55)	9 (7.04)	1.26 (0.63–2.53)	
Prothrombin G20210A	G	793 (97.66)	127 (99.21)	1.01 (0.99–1.03)	0.504
	A	19 (2.34)	1 (0.79)	0.33 (0.45–2.47)	

OR=Odds Ratio; p-values are derived from the Fisher Exact Test; CI: Confidence Interval

coronary artery disease (CAD) and control groups regarding FV G1691A genotype frequencies ($p=0.170$). Similarly, *Table I* displays the FV Gene H1299R genotype frequencies, revealing no significant difference between the control and CAD groups ($p=0.544$).

Table I also presents the genotype frequencies of FII G20210A. The results indicate no statistically significant difference between the groups regarding FII G20210A genotype frequencies ($p=0.500$). *Table II* shows the allele frequencies for the FV G1691A, FII G20210A, and FV Gene H1299R polymorphisms.

No statistically significant difference was found in allele frequencies between the patient and control groups for all three gene polymorphisms ($p>0.05$).

Discussion

Our study investigated the genotype and allele frequencies of FV G1691A (Leiden), FV H1299R, and FII G20210A variations concerning coronary artery disease (CAD) in the Turkish population. The results showed no significant differences between these mutations' genotype and allele frequencies in

the CAD and control groups. This suggests that these gene polymorphisms may not play a significant role in the development of CAD in the studied population.

Our findings are consistent with previous studies that have reported conflicting results regarding the association between these gene polymorphisms and CAD. Some studies have reported an increased risk of CAD associated with these variations, while others have found no significant association.

The prevalence of FV G1691A (Leiden) and FV H1299R variations in our study aligns with findings from studies on Turkish and Greek populations. Still, it differs from the European and Asian populations, suggesting genetic variations among different populations (22–31). These discrepancies can be attributed to genetic variations among different populations.

Similarly, the association between FII G20210A variation and CAD has yielded conflicting results in previous studies. Some studies have suggested an

increased risk (32–36), while others have found no significant association (37, 38). Our study did not find a significant difference in the genotype and allele frequencies of FII G20210A between the CAD and control groups.

In conclusion, our study found no significant association between FV G1691A (Leiden), FV Gene H1299R, and FII G20210A variations and CAD in the studied Turkish population. The prevalence of these variations aligned with studies on Turkish and Greek populations but differed from the European population. Further research with larger sample sizes and diverse populations is needed to understand these variations' role in CAD development comprehensively.

Conflict of interest statement

All the authors declare that they have no conflict of interest in this work.

References

1. Sergi C, Al Jishi T, Walker M. Factor V Leiden mutation in women with early recurrent pregnancy loss: a meta-analysis and systematic review of the causal association. *Arch Gynecol Obstet* 2015 Mar; 291(3): 671–9.
2. Van Cott EM, Khor B, Zehnder JL. Factor V Leiden. *Am J Hematol* 2016 Jan; 91(1): 46–9.
3. Kottke-Marchant K. Genetic polymorphisms associated with venous and arterial thrombosis: an overview. *Arch Pathol Lab Med* 2002; 126: 295–304.
4. Mueller T, Marschon R, Dieplinger B, Haidinger D, Gegenhuber A, Poelz W, et al. Factor V Leiden, prothrombin G20210A, and methylenetetrahydrofolate reductase C677T mutations are not associated with chronic limb ischemia: The Linz Peripheral Arterial Disease (LIPAD) Study *J Vasc Surg* 2005; 41: 808–15.
5. Kujovich J. Prothrombin-Related Thrombophilia. 2006 Jul 25 [Updated 2021 Feb 4]. In: Adam MP, Ardinger HH, Pagon RA, et al., editors. *GeneReviews®* [Internet]. Seattle (WA): University of Washington, Seattle; 1993–2022. Available at: <http://www.ncbi.nlm.nih.gov/books/NBK1148/>
6. Lunghi B, Lacoviello L, Gemmati D, Dilasio MG, Castoldi E, Pinotti M, et al. Detection of new polymorphic markers in the factor V gene: association with factor V levels in plasma. *Thromb. Haemost* 1996; 75: 45–8.
7. Castoldi E, Rosing J, Girelli D, Hoekema L, Lunghi B, Mingozzi F, et al. Mutations in the R2 FV gene affect the ratio between the two FV isoforms in plasma. *Thromb Haemost* 2000; 83: 362–5.
8. Zaatar G.S, Otrrock Z.K, Sabbagh A.S, Mahfouz R.A. Prevalence of factor V R2 (H1299R) polymorphism in the Lebanese population. *Pathology* 2006; 38: 442–4.
9. Kuhli-Hattenbach C, Hellstern P, Nögler DK, Kohnen T, Hattenbach LO, et al. Prothrombin polymorphism A19911G, factor V HR2 haplotype A4070G, and plasminogen activator-inhibitor-1 polymorphism 4G/5G and the risk of retinal vein occlusion. *Ophthalmic Genet* 2017 Sep–Oct; 38(5): 413–7.
10. Bernardi F, Faioni E.M, Castoldi E, Lunghi B, Castaman G, Sacchi E, et al. A factor V genetic component differing from factor V R506Q contributes to the activated protein C resistance phenotype. *Blood* 1997; 90: 1552–7.
11. de Visser MCH, Guasch JF, Kamphuisen PW, Vos HL, Rosendaal FR, Bertina RM. The HR2 haplotype of factor V: effects on factor V levels, normalized activated protein C sensitivity ratios and the risk of venous thrombosis. *Thromb Haemost* 2000; 83: 577–82.
12. Aleksova A, Di Nucci M, Gobbo M, Pradella P, Salam K, Barbati G, et al. Factor-V HR2 haplotype and thromboembolic disease. *Acta Cardiol* 2015 Dec; 70(6): 707–11.
13. Castoldi E, Simioni P, Kalafatis M, B Lunghi B, D Tormene D, Girelli D, et al. Combinations of 4 mutations (FV R506Q, FV H1299R, FV Y1702C, PT 20210G/A) affecting the prothrombinase complex in a thrombophilic family. *Blood* 2000; 96: 1443–8.
14. Dziadosz M, Baxi L.V. Global prevalence of prothrombin gene mutation G20210A and implications in women's health: a systematic review. *Blood Coagul Fibrinolysis* 2016; 27: 481–4.
15. Poort SR, Rosendaal FR, Reitsma PH, Bertina RM. A common genetic variation in the 3'-untranslated region of the prothrombin gene is associated with elevated plasma prothrombin levels and an increase in venous thrombosis. *Blood* 1996; 88: 3698–703.

16. Simioni P, Tormene D, Manfrin D, Gavasso S, Luni S, Stocco D, et al. Prothrombin antigen levels in symptomatic and asymptomatic carriers of the 20210A prothrombin variant. *Br J Haematol* 1998; 103: 1045–450.
17. Sabbagh AS, Ibrahim G, Kanan Z, Shammaa DMR, Khalek RA, Ghasham M, et al. Prevalence of the prothrombin G20210A polymorphism in the Lebanese population: use of a reverse hybridization strip assay approach. *Mol Biol Rep* 2007; 36: 399–403.
18. Rosendaal FR, Siscovick DS, Schwartz SM, Psaty BM, Raghunathan TE, Vos HL et al. A common prothrombin variant (20210 G to A) increases the risk of myocardial infarction in young women. *Blood* 1997; 90: 1747–450.
19. Shemesh A1, Hoffman R, Nadir Y, Keren-Politansky A, Monreal M, Brenner B, et al. Clinical significance of prothrombin G20210A mutation in homozygous patients. *Am J Hematol* 2017 Oct; 92(10): E618–E620.
20. Leroyer C, Mercier B, Oger E, Chenu E, Abgrall JF, Férec C, et al. Prevalence of 20210 A allele of the prothrombin gene in venous thromboembolism patients. *Thromb Haemost* 1998; 80: 49–51.
21. Yilmaz S, Bayan K, Tüzün Y, Batun S, A Altınta A. A comprehensive analysis of 12 thrombophilic mutations and related parameters in patients with inflammatory bowel disease: data from Turkey. *J Thromb Thrombolysis* 2006 Dec; 22(3): 205–12.
22. Rees DC, Cox M, Clegg JB. World distribution of factor V Leiden. *Lancet* 1995; 346: 1133–4.
23. Lucotte G, Mercier G. Population genetics of factor V Leiden in Europe. *Blood Cells Mol Dis* 2001; 27: 362–7.
24. Dajani R, Fatahallah R, Dajani A, Al-Shboul M, Khader Y, et al. Prevalence of coagulation factor II G20210A and factor V G1691A Leiden polymorphisms in Chechans, a genetically isolated population in Jordan. *Mol Biol Rep* 2012; 39: 9133–8.
25. Agaoglu N, Türkylmaz S, Ovalı E, Uçar F, Agaoglu C. Prevalence of prothrombotic abnormalities in patients with acute mesenteric ischemia. *World J Surg* 2005; 29: 1135–8.
26. Akar N, Akar E, Dalgin G, Sözüöz A, Ömürlü K, Cin S. Frequency of Factor V (1691 G --> A) mutation in Turkish population. *Thromb Haemost* 1997; 78: 1527–8.
27. Atasay B, Arsan S, Günlemez A, Kemahli S, Akar N. Factor V Leiden and prothrombin gene 20210A variant in neonatal thromboembolism and in healthy neonates and adults: a study in a single center. *Pediatr Hematol Oncol* 2003; 20: 627–634.
28. Eroglu Z, Biray Avcı C, Kilic M, Kosova B, Özen E, Gündüz C, et al. The prevalence of Factor V Leiden gene variant analysis of donor and recipient at the organ transplantation. *Cent. Ege Univ. Ege Tip Dergisi* 2006; 45: 185–9.
29. Gürgey A, Mesci L. The prevalence of factor V Leiden (1691 G-->A) mutation in Turkey. *Turk J Pediatr* 1997; 39: 313–5.
30. Ozbek U, Tangün Y. Frequency of factor V Leiden in Turkey. *Int J Hematol* 1996; 164: 291–2.
31. Ulukus M, Eroglu Z, Yeniel AO, Toprak E, Kosova B, Turan Ö, et al. Frequency of Factor V Leiden (Gc1691G>A), prothrombin (G20210A) and methyl-enetetrahydrofolate reductase (C677T) genes variants in women with adverse pregnancy outcome. *J. Turkish-German Gynecol Assoc* 2006; 7: 195–201.
32. Russo C, Girelli D, Olivieri O, Guarini P, Manzato F, Pizzolo F, et al. G20210A prothrombin gene polymorphism and prothrombin activity in subjects with or without angiographically documented coronary artery disease. *Circulation* 2001; 103: 2436–40.
33. Ridker PM, Hennekens CH, Miletich JP. G20210A mutation in prothrombin gene and risk of myocardial infarction, stroke, and venous thrombosis in a large cohort of US men. *Circulation* 1999; 99: 999–1004.
34. Croft SA, Daly ME, Steeds RP, KS Channer KS, Samani NJ, Hampton KK, et al. The prothrombin 20210A allele and its association with myocardial infarction. *Thromb Haemost* 1999; 81: 861–4.
35. Ardissino D, Mannucci PM, Merlini PA, Duca F, Fétiqueau R, Tagliabue L, et al. Prothrombotic genetic risk factors in young survivors of myocardial infarction. *Blood* 1999; 94: 46–51.
36. Redondo M, Watzke HH, Stucki B, Sulzer I, Biasiutti FD, Binder BR, et al. Coagulation factors II, V, and X, prothrombin gene 20210 G-.A transition, and factor V Leiden in coronary artery disease. High clotting activity is an independent risk factor for myocardial infarction. *Arterioscl Thromb Vasc Biol* 1999; 19: 1020–5.
37. Burzotta F, Paciaroni K, De Stefano V, Crea F, Maseri A, Leone G, et al. Prothrombin gene polymorphism and coronary ischaemic syndromes: a phenotype-specific meta-analysis of 12 034 subjects *Heart* 2004; 90: 82–6.
38. Ercan B, Tamer L, Sucu N, Pekdemir H, Çamsarı A, Atik U. Factor VLeiden and Prothrombin G20210A Gene Polymorphisms in Patients with Coronary Artery Disease. *Yonsei Med J* 2008; 49: 237–43.

Received: November 22, 2024

Accepted: January 14, 2025